

NAG Fortran Library Chapter Introduction**S – Approximations of Special Functions****Contents**

1	Scope of the Chapter	2
2	Background to the Problems	2
2.1	Functions of a Single Real Argument	2
2.2	Approximations to Elliptic Integrals	4
2.3	Bessel and Airy Functions of a Complex Argument	5
3	Recommendations on Choice and Use of Available Routines	5
3.1	Elliptic Integrals	5
3.2	Bessel and Airy Functions	6
4	Index	6
5	Routines Withdrawn or Scheduled for Withdrawal	8
6	References	8

1 Scope of the Chapter

This chapter is concerned with the provision of some commonly occurring physical and mathematical functions.

2 Background to the Problems

The majority of the routines in this chapter approximate real-valued functions of a single real argument, and the techniques involved are described in Section 2.1. In addition the chapter contains routines for elliptic integrals (see Section 2.2), Bessel and Airy functions of a complex argument (see Section 2.3), exponential of a complex argument, and complementary error function of a complex argument.

2.1 Functions of a Single Real Argument

Most of the routines for functions of a single real argument have been based on truncated Chebyshev expansions. This method of approximation was adopted as a compromise between the conflicting requirements of efficiency and ease of implementation on many different machine ranges. For details of the reasons behind this choice and the production and testing procedures followed in constructing this chapter see Schonfelder (1976).

Basically, if the function to be approximated is $f(x)$, then for $x \in [a, b]$ an approximation of the form

$$f(x) = g(x) \sum_{r=0}^n C_r T_r(t)$$

is used (\sum' denotes, according to the usual convention, a summation in which the first term is halved), where $g(x)$ is some suitable auxiliary function which extracts any singularities, asymptotes and, if possible, zeros of the function in the range in question and $t = t(x)$ is a mapping of the general range $[a, b]$ to the specific range $[-1, +1]$ required by the Chebyshev polynomials, $T_r(t)$. For a detailed description of the properties of the Chebyshev polynomials see Clenshaw (1962) and Fox and Parker (1968).

The essential property of these polynomials for the purposes of function approximation is that $T_n(t)$ oscillates between ± 1 and it takes its extreme values $n + 1$ times in the interval $[-1, +1]$. Therefore, provided the coefficients C_r decrease in magnitude sufficiently rapidly the error made by truncating the Chebyshev expansion after n terms is approximately given by

$$E(t) \simeq C_n T_n(t).$$

That is, the error oscillates between $\pm C_n$ and takes its extreme value $n + 1$ times in the interval in question. Now this is just the condition that the approximation be a mini-max representation, one which minimizes the maximum error. By suitable choice of the interval, $[a, b]$, the auxiliary function, $g(x)$, and the mapping of the independent variable, $t(x)$, it is almost always possible to obtain a Chebyshev expansion with rapid convergence and hence truncations that provide near mini-max polynomial approximations to the required function. The difference between the true mini-max polynomial and the truncated Chebyshev expansion is seldom sufficiently great enough to be of significance.

The evaluation of the Chebyshev expansions follows one of two methods. The first and most efficient, and hence the most commonly used, works with the equivalent simple polynomial. The second method, which is used on the few occasions when the first method proves to be unstable, is based directly on the truncated Chebyshev series, and uses backward recursion to evaluate the sum. For the first method, a suitably truncated Chebyshev expansion (truncation is chosen so that the error is less than the **machine precision**) is converted to the equivalent simple polynomial. That is, we evaluate the set of coefficients b_r such that

$$y(t) = \sum_{r=0}^{n-1} b_r t^r = \sum_{r=0}^{n-1} C_r T_r(t).$$

The polynomial can then be evaluated by the efficient Horner's method of nested multiplications,

$$y(t) = (b_0 + t(b_1 + t(b_2 + \dots t(b_{n-2} + tb_{n-1}))) \dots).$$

This method of evaluation results in efficient routines but for some expansions there is considerable loss of accuracy due to cancellation effects. In these cases the second method is used. It is well known that if

$$\begin{aligned}
b_{n-1} &= C_{n-1} \\
b_{n-2} &= 2tb_{n-1} + C_{n-2} \\
b_j &= 2tb_{j+1} - b_{j+2} + C_j, \quad j = n-3, n-4, \dots, 0
\end{aligned}$$

then

$$\sum_{r=0}^n C_r T_r(t) = \frac{1}{2}(b_0 - b_2)$$

and this is always stable. This method is most efficiently implemented by using three variables cyclically and explicitly constructing the recursion.

That is,

$$\begin{aligned}
\alpha &= C_{n-1} \\
\beta &= 2t\alpha + C_{n-2} \\
\gamma &= 2t\beta - \alpha + C_{n-3} \\
\alpha &= 2t\gamma - \beta + C_{n-4} \\
\beta &= 2t\alpha - \gamma + C_{n-5} \\
&\vdots \\
\text{say } \alpha &= 2t\gamma - \beta + C_2 \\
\beta &= 2t\alpha - \gamma + C_1 \\
y(t) &= t\beta - \alpha + \frac{1}{2}C_0
\end{aligned}$$

The auxiliary functions used are normally functions compounded of simple polynomial (usually linear) factors extracting zeros, and the primary compiler-provided functions, sin, cos, ln, exp, sqrt, which extract singularities and/or asymptotes or in some cases basic oscillatory behaviour, leaving a smooth well-behaved function to be approximated by the Chebyshev expansion which can therefore be rapidly convergent.

The mappings of $[a, b]$ to $[-1, +1]$ used range from simple linear mappings to the case when b is infinite, and considerable improvement in convergence can be obtained by use of a bilinear form of mapping. Another common form of mapping is used when the function is even; that is, it involves only even powers in its expansion. In this case an approximation over the whole interval $[-a, a]$ can be provided using a mapping $t = 2(x/a)^2 - 1$. This embodies the evenness property but the expansion in t involves all powers and hence removes the necessity of working with an expansion with half its coefficients zero.

For many of the routines an analysis of the error in principle is given, namely, if E and ∇ are the absolute errors in function and argument and ϵ and δ are the corresponding relative errors, then

$$E \simeq |f'(x)|\nabla$$

$$E \simeq |xf'(x)|\delta$$

$$\epsilon \simeq \left| \frac{xf'(x)}{f(x)} \right| \delta.$$

If we ignore errors that arise in the argument of the function by propagation of data errors, etc., and consider only those errors that result from the fact that a real number is being represented in the computer in floating-point form with finite precision, then δ is bounded and this bound is independent of the magnitude of x . For example, on an 11-digit machine

$$|\delta| \leq 10^{-11}.$$

(This of course implies that the absolute error $\nabla = x\delta$ is also bounded but the bound is now dependent on x .) However, because of this the last two relations above are probably of more interest. If possible the relative error propagation is discussed; that is, the behaviour of the error amplification factor $|xf'(x)/f(x)|$ is described, but in some cases, such as near zeros of the function which cannot be extracted explicitly, absolute error in the result is the quantity of significance and here the factor $|xf'(x)|$ is described. In general, testing of the functions has shown that their error behaviour follows fairly well these theoretical error behaviours. In regions where the error amplification factors are less than or of the order of one, the

errors are slightly larger than the above predictions. The errors are here limited largely by the finite precision of arithmetic in the machine, but ϵ is normally no more than a few times greater than the bound on δ . In regions where the amplification factors are large, of order ten or greater, the theoretical analysis gives a good measure of the accuracy obtainable.

It should be noted that the definitions and notations used for the functions in this chapter are all taken from Abramowitz and Stegun (1972). Users are strongly recommended to consult this book for details before using the routines in this chapter.

2.2 Approximations to Elliptic Integrals

Four functions provided here are symmetrised variants of the classic elliptic integrals. These alternative definitions have been suggested by Carlson (1965), Carlson (1977a) and Carlson (1977b) and he also developed the basic algorithms used in this chapter.

The standard integral of the first kind is represented by

$$R_F(x, y, z) = \frac{1}{2} \int_0^\infty \frac{dt}{\sqrt{(t+x)(t+y)(t+z)}},$$

where $x, y, z \geq 0$ and at most one may be equal to zero.

The normalization factor, $\frac{1}{2}$, is chosen so as to make

$$R_F(x, x, x) = 1/\sqrt{x}.$$

If any two of the variables are equal, R_F degenerates into the second function

$$R_C(x, y) = R_F(x, y, y) = \frac{1}{2} \int_0^\infty \frac{dt}{\sqrt{t+x}(t+y)},$$

where the argument restrictions are now $x \geq 0$ and $y \neq 0$.

This function is related to the logarithm or inverse hyperbolic functions if $0 < y < x$, and to the inverse circular functions if $0 \leq x \leq y$.

The integrals of the second kind are defined by

$$R_D(x, y, z) = \frac{3}{2} \int_0^\infty \frac{dt}{\sqrt{(t+x)(t+y)(t+z)}^3}$$

with $z > 0$, $x \geq 0$ and $y \geq 0$, but only one of x or y may be zero.

The function is a degenerate special case of the integral of the third kind

$$R_J(x, y, z, \rho) = \frac{3}{2} \int_0^\infty \frac{dt}{\sqrt{(t+x)(t+y)(t+z)(t+\rho)}}$$

with $\rho \neq 0$ and $x, y, z \geq 0$ with at most one equality holding. Thus $R_D(x, y, z) = R_J(x, y, z, z)$. The normalization of both these functions is chosen so that

$$R_D(x, x, x) = R_J(x, x, x, x) = 1/(x\sqrt{x}).$$

The algorithms used for all these functions are based on duplication theorems. These allow a recursion system to be established which constructs a new set of arguments from the old using a combination of arithmetic and geometric means. The value of the function at the original arguments can then be simply related to the value at the new arguments. These recursive reductions are used until the arguments differ from the mean by an amount small enough for a Taylor series about the mean to give sufficient accuracy when retaining terms of order less than six. Each step of the recurrences reduces the difference from the mean by a factor of four, and as the truncation error is of order six, the truncation error goes like $(4096)^{-n}$, where n is the number of iterations.

The above forms can be related to the more traditional canonical forms (see Section 17.2 in Abramowitz and Stegun (1972)).

If we write $q = \cos^2 \phi$, $r = 1 - m \cdot \sin^2 \phi$, $s = 1 + n \cdot \sin^2 \phi$, where $0 < \phi \leq \frac{1}{2}\pi$, we have

the elliptic integral of the first kind:

$$F(\phi | m) = \int_0^{\sin \phi} (1 - t^2)^{-1/2} (1 - mt^2)^{-1/2} dt = \sin \phi . R_F(q, r, 1);$$

the elliptic integral of the second kind:

$$\begin{aligned} E(\phi | m) &= \int_0^{\sin \phi} (1 - t^2)^{-1/2} (1 - mt^2)^{1/2} dt \\ &= \sin \phi . R_F(q, r, 1) - \frac{1}{3} m . \sin^3 \phi . R_D(q, r, 1) \end{aligned}$$

the elliptic integral of the third kind:

$$\begin{aligned} \Pi(n; \phi | m) &= \int_0^{\sin \phi} (1 - t^2)^{-1/2} (1 - mt^2)^{-1/2} (1 + nt^2)^{-1} dt \\ &= \sin \phi . R_F(q, r, 1) - \frac{1}{3} n . \sin^3 \phi . R_J(q, r, 1, s). \end{aligned}$$

Also the complete elliptic integral of the first kind:

$$K(m) = \int_0^{\pi/2} (1 - m . \sin^2 \theta)^{-1/2} d\theta = R_F(0, 1 - m, 1);$$

the complete elliptic integral of the second kind:

$$E(m) = \int_0^{\pi/2} (1 - m . \sin^2 \theta)^{1/2} d\theta = R_F(0, 1 - m, 1) - \frac{1}{3} m . R_D(0, 1 - m, 1).$$

2.3 Bessel and Airy Functions of a Complex Argument

The routines for Bessel and Airy functions of a real argument are based on Chebyshev expansions, as described in Section 2.1. The routines for functions of a complex argument, however, use different methods. These routines relate all functions to the modified Bessel functions $I_\nu(z)$ and $K_\nu(z)$ computed in the right-half complex plane, including their analytic continuations. I_ν and K_ν are computed by different methods according to the values of z and ν . The methods include power series, asymptotic expansions and Wronskian evaluations. The relations between functions are based on well known formulae (see Abramowitz and Stegun (1972)).

3 Recommendations on Choice and Use of Available Routines

Note: please refer to the Users' Note for your implementation to check that a routine is available.

3.1 Elliptic Integrals

IMPORTANT ADVICE: users who encounter elliptic integrals in the course of their work are strongly recommended to look at transforming their analysis directly to one of the Carlson forms, rather than to the traditional canonical Legendre forms. In general, the extra symmetry of the Carlson forms is likely to simplify the analysis, and these symmetric forms are much more stable to calculate.

The routine S21BAF for R_C is largely included as an auxiliary to the other routines for elliptic integrals. This integral essentially calculates elementary functions, e.g.,

$$\ln x = (x - 1) . R_C\left(\left(\frac{1+x}{2}\right)^2, x\right), \quad x > 0;$$

$$\arcsin x = x . R_C(1 - x^2, 1), |x| \leq 1;$$

$$\operatorname{arcsinh} x = x . R_C(1 + x^2, 1), \text{ etc.}$$

In general this method of calculating these elementary functions is not recommended as there are usually much more efficient specific routines available in the Library. However, S21BAF may be used, for example, to compute $\ln x/(x - 1)$ when x is close to 1, without the loss of significant figures that occurs when $\ln x$ and $x - 1$ are computed separately.

3.2 Bessel and Airy Functions

For computing the Bessel functions $J_\nu(x)$, $Y_\nu(x)$, $I_\nu(x)$ and $K_\nu(x)$ where x is real and $\nu = 0$ or 1 , special routines are provided, which are much faster than the more general routines that allow a complex argument and arbitrary real $\nu \geq 0$. Similarly, special routines are provided for computing the Airy functions and their derivatives $\text{Ai}(x)$, $\text{Bi}(x)$, $\text{Ai}'(x)$, $\text{Bi}'(x)$ for a real argument which are much faster than the routines for complex arguments.

4 Index

Airy function,

Ai or Ai', complex argument, optionally scaled	S17DGF
Ai, real argument	S17AGF
Ai', real argument	S17AJF
Bi or Bi', complex argument, optionally scaled	S17DHF
Bi, real argument	S17AHF
Bi', real argument	S17AKF

Arccos,

inverse circular cosine	S09ABF
-------------------------------	--------

Arccosh,

inverse hyperbolic cosine	S11ACF
---------------------------------	--------

Arcsin,

inverse circular sine	S09AAF
-----------------------------	--------

Arcsinh,

inverse hyperbolic sine	S11ABF
-------------------------------	--------

Arctanh,

inverse hyperbolic tangent	S11AAF
----------------------------------	--------

Bessel function,

J_0 , real argument	S17AEF
J_1 , real argument	S17AFF
$J_{\alpha \pm n}(z)$, real argument	S18GKF
J_ν , complex argument, optionally scaled	S17DEF
Y_0 , real argument	S17ACF
Y_1 , real argument	S17ADF
Y_ν , complex argument, optionally scaled	S17DCF

Complement of the Cumulative Normal distribution

S15ACF

Complement of the Error function,

complex argument, scaled	S15DDF
real argument	S15ADF

Cosine Integral

S13ACF

Cosine,

hyperbolic	S10ACF
------------------	--------

Cumulative Normal distribution function

S15ABF

Dawson's Integral

S15AFF

Digamma function, scaled

S14ADF

Elliptic functions, Jacobian, sn, cn, dn

complex argument	S21CBF
real argument	S21CAF

Elliptic integral,

general,	
of 2nd kind, $F(z, k', a, b)$	S21DAF
symmetrised,	
degenerate of 1st kind, R_C	S21BAF
of 1st kind, R_F	S21BBF
of 2nd kind, R_D	S21BCF
of 3rd kind, R_J	S21BDF

Erf,

real argument	S15AEF
---------------------	--------

Erfc,	
complex argument, scaled	S15DDF
real argument	S15ADF
Error function,	
real argument	S15AEF
Exponential Integral	S13AAF
Exponential,	
complex	S01EAF
Fresnel Integral,	
C	S20ADF
S	S20ACF
Gamma function	S14AAF
Gamma function,	
incomplete	S14BAF
Generalized Factorial function	S14AAF
Hankel function $H_\nu^{(1)}$ or $H_\nu^{(2)}$,	
complex argument, optionally scaled	S17DLF
Jacobian elliptic functions, sn, cn, dn,	
complex argument	S21CBF
real argument	S21CAF
Jacobian theta functions $\theta_k(x, q)$,	
real argument	S21CCF
Kelvin function,	
ber x	S19AAF
bei x	S19ABF
ker x	S19ACF
kei x	S19ADF
Legendre functions of 1st kind $P_n^m(x)$, $\overline{P}_n^m(x)$	S22AAF
Logarithm of $1 + x$	S01BAF
Logarithm of Gamma function,	
complex	S14AGF
real	S14ABF
Modified Bessel function(s),	
I_0 , real argument	S18AEF
I_1 , real argument	S18AFF
I_ν , complex argument, optionally scaled	S18DEF
K_0 , real argument	S18ACF
K_1 , real argument	S18ADF
K_ν , complex argument, optionally scaled	S18DCF
Polygamma function,	
$\psi^{(n)}(x)$, real x	S14AEF
$\psi^{(n)}(z)$, complex z	S14AFF
Psi function	S14ACF
Psi function and derivatives, scaled	S14ADF
Scaled modified Bessel function(s),	
$e^{- x }I_0(x)$, real argument	S18CEF
$e^{- x }I_1(x)$, real argument	S18CFF
$e^x K_0(x)$, real argument	S18CCF
$e^x K_1(x)$, real argument	S18CDF
Sine Integral	S13ADF
Sine,	
hyperbolic	S10ABF
Tangent,	
circular	S07AAF
hyperbolic	S10AAF
Trigamma function, scaled	S14ADF
Zeros of Bessel functions $J_\alpha(x)$, $J_\alpha'(x)$, $Y_\alpha(x)$, $Y_\alpha'(x)$	S17ALF

5 Routines Withdrawn or Scheduled for Withdrawal

None.

6 References

Abramowitz M and Stegun I A (1972) *Handbook of Mathematical Functions* (3rd Edition) Dover Publications

Carlson B C (1965) On computing elliptic integrals and functions *J. Math. Phys.* **44** 36–51

Carlson B C (1977a) Elliptic integrals of the first kind *SIAM J. Math. Anal.* **8** 231–242

Carlson B C (1977b) *Special Functions of Applied Mathematics* Academic Press

Clenshaw C W (1962) Mathematical tables *Chebyshev Series for Mathematical Functions* HMSO

Fox L and Parker I B (1968) *Chebyshev Polynomials in Numerical Analysis* Oxford University Press

Schonfelder J L (1976) The production of special function routines for a multi-machine library *Softw. Pract. Exper.* **6** (1)
